SNUTP 97-003

Quantum theory of motion of a time-dependent harmonic oscillator in the pilot-wave theory

Jeong-Young Ji* and Kwang-Sup Soh†

Department of Physics Education, Seoul National University, Seoul, 151-742, Korea

arXiv:quant-ph/9701002v1 3 Jan 1997

Abstract

The de Broglie-Bohm quantum trajectories are found in analytically closed forms for the eigenstates and the coherent state of the Lewis-Riesenfeld (LR) invariant of a time-dependent harmonic oscillator. It is also shown that an eigenstate (a coherent state) of an invariant can be interpreted as squeezed states obtained by squeezing an eigenstate (a coherent state) of another invariant. This provides ways for a whole description of squeezed states.

*Electronic address: jyji@phyb.snu.ac.kr

†Electronic address: kssoh@phyb.snu.ac.kr

The quantum theory of motion introduced by de-Broglie and Bohm provides another viewpoints on the world of quantum mechanics [1,2]. There are many examples which explain the structure of the atom, the interference and the tunneling in the quantum motional scheme [3]. The quantum trajectories have been numerically solved in a time-dependent scattering from square barriers and square potential wells [4,5]. Recently, the damped harmonic oscillator is analyzed according to de Broglie-Bohm theory [6]. However explicitly time-dependent systems have not yet been dealt with. Since the exact wave function for an explicitly time-dependent harmonic oscillator is recently found [7], we expect that one may calculate the quantum trajectories for an eigenstate, a coherent state, and a squeezed state of the system.

In this Letter, we find the de Broglie-Bohm quantum trajectories for a time-dependent harmonic oscillator. The LR invariants [8] are constructed using the creation and the annihilation operators which are expressed in terms of a classical solution. It is argued that since any linear combination of two independent classical solutions is also a solution, there are many invariants and the invariant can be specified uniquely (except for a constant factor) if we fix the classical solution. It is shown that the invariants with different solutions are obtained by squeezing each other with a squeezing operator. Here the squeezing operator describes the Bogoliubov transformation between two sets of creation and annihilation operators. For an invariant, we consider the eigenstates and coherent states as a guiding wave in the de Broglie-Bohm theory and find the corresponding quantum trajectories. Because these states are squeezed states for another invariant, we have a whole description of squeezed states. The solutions are applied to a time-independent oscillator and a damped oscillator. In the damped oscillator we find the new type of quantum trajectories which have not been found in Ref. [6]. Here the quantum motions are oscillating while damping.

In the causal interpretation of quantum mechanics, the wave is mathematically described by $\Psi(\mathbf{x}, t)$, a solution to Schrödinger's wave equation

$$i\hbar\frac{\partial\Psi}{\partial t} = \left(-\frac{\hbar^2}{2m}\nabla^2 + V\right)\Psi \tag{1}$$

Setting $\Psi = Re^{iS/\hbar}$ where R and S are real functions of position and time, the particle trajectories of a particle of mass m moving in the potential $V + Q$ is described by the following quantum Hamilton-Jacobi equation:

$$\frac{\partial S}{\partial t} + \frac{(\nabla S)^2}{2m} + V + Q = 0 \tag{2}$$

where the quantum potential is given by

$$Q = -\frac{\hbar^2}{2m}\frac{\nabla^2 R}{R} \tag{3}$$

Thus the particle trajectories associated with a given quantum mechanical state can be obtained by solving the quantum equation of motion

$$\dot{\mathbf{x}} = (1/m)\nabla S(\mathbf{x}, t)|_{\mathbf{x}=\mathbf{x}(t)} \tag{4}$$

or equivalently

$$m\ddot{\mathbf{x}} = -\nabla(V + Q)|_{\mathbf{x}=\mathbf{x}(t)}. \tag{5}$$

The LR invariant, $I(t)$, of a time-dependent harmonic oscillator

$$\hat{H}(t) = \frac{1}{2M(t)}\hat{p}^2 + \frac{1}{2}M(t)\omega^2(t)\hat{q}^2 \tag{6}$$

can be written as (hereafter we adopt $\hbar = 1$ units.)

$$\hat{I}(t) = \omega_I\left[\hat{b}^\dagger(t)\hat{b}(t) + \frac{1}{2}\right]. \tag{7}$$

Here we have used the creation and annihilation operators introduced in Ref. [9]. Noting that

$$f(t) = \sqrt{\frac{g_-(t)}{2\omega_I}}e^{-i\Theta(t,t_0)} \tag{8}$$

with $\Theta(t, t_0) = \int_{t_0}^{t} dt\frac{\omega_I}{M(t)g_-(t)}$ satisfies the classical equation of motion:

$$\frac{d}{dt}[M(t)\frac{d}{dt}f(t)] + M(t)\omega^2(t)f(t) = 0. \tag{9}$$

and using

$$\frac{d}{dt}\left[\sqrt{\frac{g_-(t)}{2\omega_I}}e^{-i\Theta(t,t_0)}\right] = -\frac{i}{M}\left[\sqrt{\frac{\omega_I}{2g_-(t)}} - i\frac{g_0(t)}{\sqrt{2\omega_I g_-(t)}}\right]e^{-i\Theta(t,t_0)} \tag{10}$$

the operators $\hat{b}$ and $\hat{b}^\dagger$ in $\hat{b}(t) = be^{-i\Theta(t,t_0)}$ and $b^\dagger(t) = b^\dagger e^{i\Theta(t,t_0)}$ can be written as

$$\begin{aligned}\hat{b} &= i[-M(t)\dot{f}^*(t)\hat{q}(t) + f^*(t)\hat{p}(t)] \\ \hat{b}^\dagger &= -i[-M(t)\dot{f}(t)\hat{q}(t) + f(t)\hat{p}(t)]\end{aligned} \tag{11}$$

where $\Theta(t,t_0) = \int_{t_0}^{t}\frac{dt}{2M(t)|f(t)|^2}$ and we normalize $f(t)$ so that $[b,b^\dagger] = iM(t)[\dot{f}f^* - f\dot{f}^*](t) = 1$. (For this condition, the Wronskian of f and f^* should not vanish, i.e. they are linearly independent.) There are uncountable number of invariants since we have freedom in choosing the classical solution $f(t)$: Any linear combination of two independent classical solutions is also a solution, or equivalently, we are free to choose the parameters c_1, c_2, and c_3 with fixed two independent solutions $f_1(t)$ and $f_2(t)$ in Eq. (7) of Ref. [9]. Hereafter when we construct the LR invariant, we choose the function $f(t)$ and ω_I instead of choosing $c_{1,2,3}$. Then the auxiliary function is $g_-(t) = 2\omega_I|f(t)|^2$ and the annihilation operator and the eigenstates of the invarint are uniquely specified only by $f(t)$. Thus we denote them by b_f and $|n\rangle_f$, respectively. With f fixed, the invariants, I_f, may differ by a constant factor ω_I, which is irrelevant to the construction of quantum states.

Let us take another classical solution, $F(t)$, which are related by $f(t)$ as

$$F(t) = uf(t) + v^*f^*(t) \tag{12}$$

$$F^*(t) = vf(t) + u^*f^*(t) \tag{13}$$

to construct another creation and annihilation operators

$$\begin{aligned}\hat{b}_F &= i[-M(t)\dot{F}^*(t)\hat{q}(t) + F^*(t)\hat{p}(t)] \\ \hat{b}_F^\dagger &= -i[-M(t)\dot{F}(t)\hat{q}(t) + F(t)\hat{p}(t)]\end{aligned} \tag{14}$$

Then the relation between two sets of creation and annihilation operators is given by the following Bogoliubov transformation or unitary (similarity) transformation:

$$\hat{b}_F = u\hat{b}_f + v^*\hat{b}_f^\dagger = \hat{S}^\dagger \hat{b}_f \hat{S} \tag{15}$$

generated by the squeezing operator

$$\hat{S} = e^{i\theta_u \hat{b}_f^\dagger \hat{b}_f} \exp\left(\frac{\sigma}{2} e^{+i(\theta_v - \theta_u)} \hat{b}_f^{\dagger 2} - \frac{\sigma}{2} e^{-i(\theta_v - \theta_u)} \hat{b}_f^2\right) \tag{16}$$

where $\sigma = \cosh^{-1}|u|$. Therefore it is verified that the eigenstate of I_F is obtained by squeezing the eigenstate of I_f as

$$|n\rangle_F = \hat{S}^\dagger |n\rangle_f \tag{17}$$

and for the coherent state $|\alpha\rangle = e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle$

$$|\alpha\rangle_F = \hat{S}^\dagger |\alpha\rangle_f . \tag{18}$$

Since $|\alpha\rangle_f$ can be obtained by displacing $|0\rangle_f$, $|\alpha\rangle_F$ is obtained by displacing and squeezing $|0\rangle_f$, i.e.

$$|\alpha\rangle_F = |\sigma, \alpha\rangle_f = \hat{S}^\dagger(\sigma)\hat{D}(\alpha)|0\rangle_f \tag{19}$$

with $\hat{D}(\alpha) = e^{\alpha\hat{a}^\dagger - \alpha^*\hat{a}}$.

Recently, in Ref. [7], we have found the explicit form of wave function for the n-th eigenstate of the invariant (7):

$$\psi_n(q,t) = R_n(q,t) e^{iS_n(q,t)/\hbar} \tag{20}$$

where

$$R_n(q,t) = \frac{1}{\sqrt{2^n n!}} \left(\frac{\omega_I}{\pi g_-(t)}\right)^{\frac{1}{4}} e^{-\frac{\omega_I}{2g_-(t)}q^2} H_n\left(\sqrt{\frac{\omega_I}{g_-(t)}} q\right). \tag{21}$$

and

$$S_n(q,t) = -\frac{g_0(t)}{2g_-(t)} q^2 - \left(n + \frac{1}{2}\right)\Theta(t, t_0). \tag{22}$$

When the exact form of the wave function is given, it is straightforward to find the quantum trajectory by solving the quantum equation of motion (4). The quantum equation of motion for the n-th eigenstate of the invariant is

$$M(t)\dot{q} = -\frac{g_0(t)}{g_-(t)}q. \tag{23}$$

Using $g_0 = -\frac{M}{2}\dot{g}_-$, we find a solution easily

$$q(t) = \frac{q(t_0)}{\sqrt{g_-(t_0)}}\sqrt{g_-(t)}. \tag{24}$$

As expected from the result of damped oscillator [6], the quantum trajectory is independent of the quantum number n in a general time-dependent oscillator.

Now we consider the coherent state of the invariant

$$\Psi_\alpha(q,t) = e^{-|\alpha|^2/2}\sum_n \frac{\alpha^n}{\sqrt{n!}}\psi_n(q,t) \tag{25}$$

Using $e^{-z^2+2xz} = \sum_0^\infty \frac{z^n}{n!}H_n(x)$ with $z = \frac{1}{\sqrt{2}}\alpha e^{-i\Theta(t,t_0)}$ and $x = \sqrt{\frac{\omega_I}{g_-}}q$, we have

$$\Psi_\alpha(q,t) = \left(\frac{\omega_I}{\pi g_-(t)}\right)^{\frac{1}{4}} e^{-\frac{|\alpha|^2}{2}-\frac{\omega_I}{2g_-(t)}q^2-i\frac{g_0}{2g_-}q^2-\frac{i}{2}\Theta(t,t_0)}e^{-z^2+2xz}. \tag{26}$$

We rewrite the above wave function as

$$\Psi_\alpha(q,t) = R_\alpha(q,t)e^{iS_\alpha(q,t)/\hbar} \tag{27}$$

where

$$R_\alpha(q,t) = \left(\frac{\omega_I}{\pi g_-(t)}\right)^{\frac{1}{4}} e^{-\frac{|\alpha|^2}{2}-\frac{\omega_I}{2g_-(t)}q^2-\frac{\alpha^2}{2}\cos[2\Theta(t,t_0)]+\alpha q\sqrt{\frac{2\omega_I}{g_-(t)}}\cos\Theta(t,t_0)} \tag{28}$$

and

$$S_\alpha(q,t) = -\frac{g_0}{2g_-}q^2 - \frac{1}{2}\Theta(t,t_0) + \frac{\alpha^2}{2}\sin[2\Theta(t,t_0)] - \alpha q\sqrt{\frac{2\omega_I}{g_-(t)}}\sin[\Theta(t,t_0)]. \tag{29}$$

Therefore, from (4), we have the quantum equation of motion for the coherent state

$$M\dot{q} = -\frac{g_0}{g_-}q - \alpha\sqrt{\frac{2\omega_I}{g_-}}\sin\Theta \tag{30}$$

which can be easily integrated using

$$\frac{d}{dt}\left(\frac{q}{\sqrt{g_-}}\right) = \alpha\sqrt{\frac{2}{\omega_I}}\frac{d}{dt}\cos\Theta, \tag{31}$$

then we have the following quantum trajectory

$$q(t) = \frac{q(t_0)}{\sqrt{g_-(t_0)}}\sqrt{g_-(t)} + \alpha\sqrt{\frac{2g_-(t)}{\omega_I}}[\cos\Theta(t,t_0) - 1]. \tag{32}$$

One should note that the coherent state (25) of an invariant can be interpreted as the displaced squeezed state of another invariant [see (19) and related remarks]. Thus the quantum motion we have found can be, in general, gives a whole description of the motion for a squeezed state.

As a simple example we consider the time-independent case with $M(t) = m$ and $\omega(t) = \omega_0$. If we choose the classical solution $f(t) = (1/\sqrt{2m\omega_0})e^{-i\omega_0 t}$ and $\omega_I = \omega_0$ we have

$$\hat{I}_f(t) = \hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega_0^2\hat{q}^2. \tag{33}$$

For an eigenstate of this invariant, it follows from $g_-(t) = 1/m$ and (24) that

$$q(t) = q(t_0), \tag{34}$$

and for a coherent state, from (32)

$$q(t) = q(0) + \alpha\sqrt{\frac{2}{m\omega_0}}(\cos\omega_0 t - 1), \tag{35}$$

which are well known results [3].

While if we choose the classical solutions as $F(t) = (\cosh\sigma)f(t) + (\sinh\sigma)f^*(t)$, we have

$$\hat{I}_F = g_-\frac{\hat{p}^2}{2} + g_0\frac{\hat{p}\hat{q} + \hat{q}\hat{p}}{2} + g_+\frac{\hat{q}^2}{2} \tag{36}$$

where

$$\begin{aligned} g_-(t) &= (1/m)(e^{2\sigma}\cos^2\omega_0 t + e^{-2\sigma}\sin^2\omega_0 t), \\ g_0\ (t) &= -(e^{2\sigma} - e^{-2\sigma})\omega_0\cos\omega_0 t\sin\omega_0 t, \\ g_+(t) &= m\omega_0^2(e^{2\sigma}\sin^2\omega_0 t + e^{-2\sigma}\cos^2\omega_0 t). \end{aligned} \tag{37}$$

and we have a quantum trajectory, for an eigenstate $|n\rangle_F$

$$q(t) = q(0)\sqrt{\cos^2\omega_0 t + e^{-4\sigma}\sin^2\omega_0 t}, \tag{38}$$

and for a coherent state $|\alpha\rangle_F$

$$q(t) = \sqrt{\cos^2\omega_0 t + e^{-4\sigma}\sin^2\omega_0 t}\left[q(0) + \alpha e^{\sigma}\sqrt{2/m\omega_0}(\cos\Theta - 1)\right], \tag{39}$$

with $\Theta(t) = \tan^{-1}[e^{-2\sigma}\tan(\omega_0 t - r\pi)] + r\pi$, for $|\omega_0 t - r\pi| \leq \pi/2$ $(r = 0, \pm 1, \pm 2, ...)$. As stated before, $|\alpha\rangle_F$ is a displaced squeezed state which is obtained by displacing and squeezing the ground state of the time-independent harmonic oscillator. Thus we have found the quantum motion for the displaced squeezed state.

For comparison with the result of Ref. [6], we take the damped oscillator as another example: $M(t) = me^{2\gamma t}$, $\omega(t) = \omega_0$ with $\omega_0 > \gamma$. The classical equation of motion is

$$\ddot{f} + 2\gamma\dot{f} + \omega_0^2 f = 0 \tag{40}$$

and if we choose $f(t) = (1/\sqrt{2m\Omega})e^{-\gamma t - i\Omega t}$ with $\Omega = \sqrt{\omega_0^2 - \gamma^2}$ as a solution and $\omega_I = \Omega$, we have $g_-(t) = (1/m)e^{-2\gamma t}$ and the quantum trajectory for $|n\rangle_f$ is

$$q(t) = q(0)e^{-\gamma t} \tag{41}$$

which is the same result obtained in Ref. [6]. However if we choose $F(t) = (\cosh\sigma)f(t) + (\sinh\sigma)f^*(t)$ we have $g_-(t) = (e^{-2\gamma t}/m)(e^{2\sigma}\cos^2\Omega t + e^{-2\sigma}\sin^2\Omega t)$, and the quantum trajectory for $|n\rangle_F$

$$q(t) = q(0)e^{-2\gamma t}(\cos^2\Omega t + e^{-4\sigma}\sin^2\Omega t). \tag{42}$$

This quantum trajectory is clearly oscillatory while damping.

In summary, we have constructed LR invariants introducing the creation and the annihilation operators which are specified by a classical solution. We have found the quantum trajectory for the eigenstates and for the coherent state of this invariant. As in the cases of LR invariant, the time evolution of quantum operator $\hat{q}(t)$ and $\hat{p}(t)$ in the Heisenberg picture, and the wave function of the invariant, the quantum trajectories are also expressed only in terms of the classical solution of the oscillator. The solutions were verified in the time-independent oscillator and a damped oscillator.

The quantum trajectory we have found can be easily applied to other time-dependent oscillators as far as the classical solutions are known. We expect that the oscillator's trajectory give a good description, as an example, of explicitly time-dependent system in the de Broglie-Bohm quantum theory. Our formalism can also be applied to a Paul trap [10] or a modified Paul trap [11] to study the quantum motion of a particle.

This work was supported by the Center for Theoretical Physics (S.N.U.) and the Basic Science Research Institute Program, Ministry of Education Project No. BSRI-96-2418. One of us (JYJ) is supported by Ministry of Education for the post-doctorial fellowship.

REFERENCES

[1] L. de Broglie, C. R. Acad. Sci. 183 (1926) 447; 184 (1927) 273; 185 (1927) 380.

[2] D. Bohm, Phys. Rev. 85 (1952) 166, 180.

[3] P.R. Holland, The quantum theory of motion (Cambridge Univ. Press, Cambridge, 1993).

[4] C. Dewdney and B. Hiley, Found. Phys. 12 (1982) 27.

[5] J. Hirschfelder and A. Christoph, Chem. Phys. 61 (1974) 5435.

[6] M.A. Vandyck, J. Phys. A 27 (1994) 1743.

[7] J.Y. Ji, J.K. Kim, S.P. Kim and K.S. Soh, Phys. Rev. A 52 (1995) 3352.

[8] H.R. Lewis Jr. and W.B. Riesenfeld, J. Math. Phys. 10 (1969) 1458.

[9] J.Y. Ji, J.K. Kim and S.P. Kim, Phys. Rev. A 51 (1995) 4268.

[10] W. Paul, Rev. Mod. Phys, 62 (1990) 531.

[11] J.C. Park, J.Y. Ji and K.S. Soh, J. Korean Phys. Soc. (to be published).